

INSTITUUT VOOR PLANTENZIEKTENKUNDIG ONDERZOEK
WAGENINGEN, NEDERLAND
DIRECTEUR: Dr. J. G. TEN HOUTEN

MEDEDELINGEN No 178 EN 179

INVESTIGATIONS ON A LEAFHOPPER-BORNE CLOVER VIRUS

DOOR
H. H. EVENHUIS


**DE VECTOREN VAN HET BLOEMVERGROENINGSVIRUS
VAN KLAVER**

**(THE VECTORS OF THE VIRUS CAUSING PHYLLODY (VIRESCENCE)
IN CLOVER FLOWERS)**


DOOR
H. H. EVENHUIS



*Done with
Conference Rep.*



OVERDRUK UIT:
PROC. 3RD CONF. ON POTATO VIRUS DISEASES: 251-254, 1958 EN
TIJDSCHRIFT OVER PLANTEZIEKTEN, 64: 335-336, 1958



INSTITUUT VOOR PLANTENZIEKTENKUNDIG ONDERZOEK (I.P.O.)

Office and main laboratory:

Binnenhaven 4a, tel. 2151, 2152 en 3641
Wageningen, The Netherlands.

Staff:

Director:

Dr. J. G. TEN HOUTEN.

Head of the Entomological Dept.:

Dr. H. J. DE FLUITER, Wageningen.

Deputy head of the Mycological Dept.:

Ir. A. M. VAN DOORN, Wageningen.

Head of the Nematological Dept.:

Dr. Ir. J. W. SEINHORST, Wageningen.

Head of the Plant Disease Resistance Dept.:

Dr. J. C. s'JACOB, Wageningen.

Head of the Virological Dept.:

Miss Dra. F. QUAK, Wageningen.

Deputy head of the Dept. for economic use of pesticides and aerial spraying in agriculture:

Miss M. C. KERSEN, Wageningen.

Research workers at the Wageningen Laboratory:

Ir. A. B. R. BEEMSTER, Virologist

Ir. R. E. LABRUYÈRE, Phytopathologist

Ir. J. A. DE BOKX, Virologist

Drs. H. P. MAAS GEESTERANUS, Phytopathologist

Dr. Ir. L. BOS, Virologist

Dr. J. C. MOOI, Phytopathologist

Ir. A. M. VAN DOORN, Phytopathologist

Ir. H. DEN OUDEN, Nematologist

Dr. H. H. EVENHUIS, Entomologist

Miss Dra. H. J. PFAELTZER, Virologist

Dr. H. J. DE FLUITER, Entomologist

Miss Dra. F. QUAK, Virologist

Dr. C. J. H. FRANSSEN, Entomologist

Dr. Ir. J. W. SEINHORST, Nematologist

Dr. J. GROSJEAN, Phytopathologist

Ir. J. VAN DER SPEK, Phytopathologist

Ir. N. HUBBELING, Phytopathologist and plantbreeder

Ir. F. H. F. G. SPIERINGS, Plantphysiologist

Dr. J. C. s'JACOB, Phytopathologist and plantbreeder

Dr. F. TJALLINGH, Phytopathologist

Dr. J. H. VENEKAMP, Biochemist

Drs. J. C. ZADOKS, Phytopathologist

Miss Dr. C. H. KLINKENBERG, Nematologist

Research workers elsewhere:

Ir. H. A. VAN HOOFF, Phytopathologist } detached to „Proefstation voor de Groenteteelt

Drs. L. E. VAN 'T SANT, Entomologist } in de volle grond", Alkmaar, tel. K 2200-4568.

Drs. D. J. DE JONG, Entomologist } detached to „Proefstation voor de Fruitteelt in de

Ir. G. S. ROOSJE, Phytopathologist } volle grond", Wilhelminadorp, Goes, tel. K 1100-2261

Ir. T. W. LEFERING, Phytopathologist/Virologist, detached to „Proeftuin Noord Limburg"
Venlo, tel. K 4700-2503.

Drs. G. SCHOLTEN, Phytopathologist, detached to „Proefstation voor de bloemisterij in Nederland", Aalsmeer, tel. K 2977-688.

Ir. G. P. TERMOHLEN, Phytopathologist, detached to „Proeftuin voor de Groente- en Fruitteelt onder glas", Naaldwijk, tel. K 1740-4545.

Guest workers:

Dr. P. A. VAN DER LAAN, Entomologist, „Laboratorium voor toegepaste Entomologie der Gemeente Universiteit," Amsterdam, tel. K 2900-56282.

Dr. Ir. G. S. VAN MARLE, Entomologist, Diepenveenseweg 226, Deventer. tel. K 6700-3617.

Ir. G. W. ANKERSMIT, Entomologist, „Laboratorium voor Entomologie", Agricultural University, Wageningen, tel. K 8370-2438.

Dr. Ir. J. B. M. VAN DINTHER, Entomologist, „Laboratorium voor Entomologie", Agricultural University, Wageningen, tel. K 8370-2438.

Aphidological Adviser:

Mr. D. HILLE RIS LAMBERS, Entomologist, T.N.O., Bennekom, tel. K 8379-458.

INVESTIGATIONS ON A LEAFHOPPER-BORNE CLOVER VIRUS

H. H. EVENHUIS

Institute of Phytopathological Research, Wageningen, the Netherlands

Some years ago MARAMOROSCH (1953) described a new leafhopper-borne virus disease from the Netherlands. He collected some leafhopper species in the vegetation of a cherry orchard at Gronsvelt in the neighbourhood of Maastricht and transferred them in the laboratory at Wageningen to different test plants.

Several specimens of *Euscelis plebejus*, if transferred to crimson clover seedlings, caused symptoms, which MARAMOROSCH attributed to a virus disease, new for Western Europe. He described two groups of symptoms: 1. the first signs, which were characterized by enlarged veins and deep ridges on the upper surface of the leaflets, often with pronounced enations on the lower surface on the veins; 2. the later symptoms, a chlorosis which progressed from the outer edges towards the midribs of the leaflets and small malformed leaves on shortened petioles.

We did some experiments in the summer of 1953 in order to get more information about the supposed virus of MARAMOROSCH.

On each of 35 crimson clover plants two to ten specimens of *Euscelis plebejus* were caged for a week. The insects were obtained from the same orchard where MARAMOROSCH collected his leafhoppers. After some time eight plants showed only the first named group of symptoms, three only the second group and three both groups of symptoms. These data show, that the two groups of symptoms may appear separately and that there is no relation between these two groups of symptoms. In the following the two groups of symptoms will be considered separately.

1. The enlarged veins, the deep ridges and the enations may already appear after a rather short time of feeding of the insect; according to MARAMOROSCH in some cases the symptoms already appeared within a week. We noticed the symptoms after a somewhat longer interval, but in most cases they were very conspicuous after two weeks. On some plants the symptoms were much more pronounced than on others; also it was noted that the same leafhoppers sometimes did produce the symptoms on one plant but not on another. Besides, no correlation could be found between the number, the size of the leafhoppers in the different instars or the size of the crimson clover plants and the intensity of the symptoms. Sometimes only one small nymph already sufficed to produce very severe symptoms. In the winter season there were mostly no symptoms at all.

The symptoms could also be produced on *Trifolium hybridum* and exceptionally also on white clover. No symptoms were obtained on red clover, pea, bean, alfalfa and white sweet clover. All further experiments have been done with crimson clover.

Besides *Euscelis plebejus* also some other leafhopper species have been tested. The taxonomy of the genus *Euscelis* is very difficult, but the work of WAGNER, RIBAUT and especially the recent biological studies of MÜLLER (1954, 1955) have contributed considerably to our knowledge on this subject. The taxonomy has not yet been entirely elucidated, but it appears that practically all forms, found regularly in the grass, belong to the species *E. plebejus* and *E. lineolatus* and both species are equally capable of

producing the symptoms on crimson clover. Both species are very polyphagous and may breed on widely different plant species.

No symptoms were produced on crimson clover by *Aphrodes bicinctus*, *Macrosteles spec.* and *Arthaleus pascuellus*. The latter is a grass feeding species that cannot live for a long time on crimson clover.

Since symptoms as the above mentioned can be caused by a virus as well as by a toxic substance, the nature of the causative agent should be determined experimentally. If it would be possible to rear a strain of *Euscelis*, which does not produce symptoms on healthy clover plants until after a period of feeding on diseased plants, proof of the virus nature of the agent would be delivered.

Attempts to produce such a strain have, however, not been successful, not even if insects were hatched from isolated eggs between wet filterpaper in Petri dishes. Nymphs reared this way were capable of producing the symptoms on healthy plants. From this fact we must conclude that if a virus is concerned it must have been transmitted easily through the egg.

An argument against the virus nature of the agent, however, is the fact that if a leafhopper is put on a healthy plant, the symptoms occur in the leaves unfolding subsequently, but not in those unfolding some time after the insect has been removed from the plant. For instance: on a plant on which five leafhoppers had been feeding from September 11 to 14, distinct symptoms appeared only on one leaf on September 26, the younger leaves showing no symptoms at all. On another plant a nymph had been feeding from May 28 till June 24. Distinct symptoms appeared between June 17 and July 15 on seven leaves, whereas on the younger leaves no symptoms appeared.

If one puts a little cage with a leafhopper round a leafstalk, the new, unfolding leaves show the symptoms. This implies that the symptoms are induced by a toxic substance that spreads through the plant, through the leaf stalk to the vegetation point.

In literature more cases are known of leafhoppers producing viruslike symptoms, caused by a systemic toxic substance. KUNKEL (1933) describes a wilting of young peach seedlings caused by *Macropsis trimaculata*; SEVERIN, DOUGLAS HORN & FRAZIER (1945) describe viruslike symptoms on sugar beet and aster induced by the feeding of the leafhopper *Xerophloea vanduzeei*; SEVERIN (1947) in California mentions the appearance of symptoms by feeding of specimens of 10 leafhopper vectors of aster yellows on aster and celery; BENNETT (1952) reports a vein-clearing on sugar beet caused by a systemic toxic substance, introduced into the plant by the feeding of some specimens of *Circulifer tenellus*.

2. The second group of symptoms, described by MARAMOROSCH, may appear after a much longer period; in most cases not until after five or six weeks. From 1953 until 1956 we were not able to reproduce these symptoms. However, in 1956 we found at Gronsvelt, near Maastricht, a red clover plant with viruslike symptoms. These symptoms appeared to be produced by a virus, that was transmitted by the leafhoppers *Euscelis plebejus* and *E. lineolatus* from diseased plants to healthy red clover seedlings. Leafhoppers, reared in cages on healthy seedlings, did not produce these symptoms.

The symptoms were a distinct vein-chlorosis or vein-clearing on the leaves, which seemed darker than the healthy ones; later on a stunting and proliferation of the plant, a yellowing of the young leaves beginning at the edges, a production of very small leaves and a phyllody of the flowers appeared. The symptoms proved to be identical with those, described by FRAZIER & POSNETTE (1956), who were the first to prove that this disease is really a virus disease, transmitted by leafhoppers.

The virus was also transmitted by *Euscelis plebejus* and *E. lineolatus* to crimson clover, producing the same symptoms. So we arrived at the conclusion that the second group of symptoms on crimson clover, described by MARAMOROSCH, are produced by the same virus. However, in 1953 a phyllody of the flowers was never observed, because of the fact that the plants with virus symptoms did not flower at all. We were also able to prove, that *Aphrodes bicinctus* is capable to transmit the virus. In the meantime FRAZIER & POSNETTE (1957) found that the virus they described in 1956 is in fact a complex of two viruses of which one only produces the phyllody and the other especially the symptoms of witches' broom.

Up till now we were not able to separate the two viruses, so when speaking about the clover virus it may be possible that two viruses are concerned.

We were able to transmit the virus by means of *Euscelis spec.* in one of four cases to aster. On this aster plant we got a slight chlorosis of the leaves, but a very pronounced phyllody with green, rather small flower heads and especially phyllody of the ovary, just as in clover. From the symptoms on aster we conclude, that the virus disease of clover is not identical with the european aster yellows, described by HENZE & KUNZE (1955), nor with the American aster yellows (eastern and western strain).

The symptoms, shown by red and crimson clover, were also obtained on white clover. By feeding virus-infected leafhoppers on peas also a phyllody was obtained. We also used *Cuscuta subinclusa* in our experiments; by means of this dodder species the virus could easily be transmitted from red clover to red clover.

No exact data are known about the period, necessary for the leafhoppers to pick up the virus from a diseased plant and to transmit it to a healthy plant. At any rate four days feeding on a diseased plant and three days feeding on a healthy plant are sufficient. Probably much shorter times would also be sufficient to obtain virus transmission.

The latent period in the vector is very long. By transferring every week leafhoppers that had been feeding on a diseased plant to a new healthy seedling of red clover in the glasshouse, the latent period in the vector was estimated to amount six to seven weeks or even more.

The nymphs as well as the adults can pick up the virus. As the latent period lasts very long, the nymphs had become adults when the insects became infective.

Phyllody of clover in Western Europe is already known for a very long time. It was already mentioned in the handbook of MASTERS (1869). Also PENZIG (1890) states, that phyllody of white clover is very common. In earlier times it was ascribed to several different causes, which BECKER (1941) mentions in his paper. FRAZIER & POSNETTE (1956) were the first to prove that a phyllody of clover is caused by a virus, transmitted by leafhoppers. They were able to transmit the virus by dodder to strawberries on which the typical symptoms of the green petal disease of strawberries developed.

So the symptoms, described by MARAMOROSCH from crimson clover and caused by feeding of the leafhopper *Euscelis spec.* may be divided in two distinct groups: 1. symptoms which are not caused by a virus, but by a toxic substance, spreading systemically through the plant; 2. symptoms, caused by a virus, identical with the virus, described by FRAZIER & POSNETTE. It can be transmitted by some leafhopper species to a number of plant species.

The first group of symptoms have only been observed, after *Euscelis spec.* had been feeding on crimson clover, on *Trifolium hybridum* and in one case on white clover.

SUMMARY

A further study was made of the leafhopper-borne disease of crimson clover described from the Netherlands by MARAMOROSCH. The first symptoms, principally consisting of deep ridges on the upper side of the leaflets, proved to be caused by a toxic substance excreted by the leafhopper. These symptoms appeared rather shortly (sometimes already after one week) after the leafhoppers had fed on the plant. The later symptoms which were among others a chlorosis, progressing from the margins towards the midribs of the leaflets, appeared after a much longer period (minimum: 3 to 4 weeks). These symptoms proved to be caused by a virus, transmitted by *Euscelis plebejus*, *E. lineolatus*, and also by *Aphrodes bicinctus*. The virus involved is probably the same as the one causing green petal disease of strawberry, described by FRAZIER & POSNETTE. Judged by the symptoms produced in aster the virus is not identical with the virus causing European aster yellows, described by HEINZE & KUNZE, or with the eastern and western strains of aster yellows in the United States of America.

REFERENCES

- BECKER, R., - 1941. Untersuchungen über die Blüh- und Befruchtungsverhältnisse und die Blütenmissbildungen des Weissklee (*Trifolium repens* L.). *Futtersaatbau* 1: 1-84.
- BENNET, C. W., - 1952. Vein clearing on sugar beet induced by the beet leafhopper. *Phytopath.* 42: 535-537.
- FRAZIER, N. W. & A. F. POSNETTE, - 1956. Leafhopper transmission of a clover virus causing green petal disease in strawberry. *Nature* 177: 1040-1041.
- FRAZIER, N. W. & A. F. POSNETTE, - 1957. Transmission and host-range studies of strawberry green-petal virus. *Ann. appl. Biol.* 45: 580-588.
- HEINZE, K. & L. KUNZE, - 1955. Die europäische Asterolgelbsucht und ihre Übertragung durch Zwergzikaden. *Nachr.-Bl. dtsh. Pfl.-Sch.-D. (Braunsch.)* 7: 161-164.
- KUNKEL, L. O., - 1933. Insect transmission of peach yellows. *Contr. Boyce Thomps. Inst.* 5: 19-28.
- MARAMOROSCH, K., - 1953. A new leafhopper-borne plant disease from Western Europe. *Pl. Dis. Rep.* 37: 612-613.
- MASTERS, M. T., - 1869. *Vegetable teratology*. London.
- MÜLLER, H. J., - 1954. Der Saisondimorphismus bei Zikaden der Gattung *Euscelis* Brullé (*Homoptera Auchenorrhyncha*) (3. Beitrag zur Biologie mitteleuropäischer Zikaden). *Beitr. Entom.* 4: 1-56.
- MÜLLER, H. J., - 1955. Die Bedeutung der Tageslänge für die Saisonformbildung der Insekten, insbesondere bei den Zikaden. *Ber. 7 Wanderversamml. dtsh. Entom.*: 102-120.
- PENZIG, M., - 1890. *Pflanzen-Teratologie*, 1. Band. Genua.
- SEVERIN, H. H. P., - 1947. Plant symptoms induced by feeding of some leafhopper species. *Hilgardia* 17: 219-221.
- SEVERIN, H. H. P., F. DOUGLAS HORN & N. W. FRAZIER, - 1945. Certain symptoms resembling those of curly top or aster yellows, induced by saliva of *Xerophloea vanduzeei*. *Hilgardia* 16: 337-350.

DE VECTOREN VAN HET BLOEMVERGROENINGSVIRUS VAN KLAVER ¹⁾

*With a summary: The vectors of the virus causing phyllody (virescence)
in clover flowers*

DOOR

H. H. EVENHUIS

Instituut voor Plantenziektenkundig Onderzoek, Wageningen

Bloemvergroeningsverschijnselen zijn al minstens sedert de vorige eeuw uit West-Europa bekend (Bos, 1957). Door FRAZIER & POSNETTE (1956) is voor het eerst aangetoond, dat deze verschijnselen kunnen worden veroorzaakt door een virus, dat door cicaden wordt overgebracht. In 1957 vermeldde de beide genoemde auteurs, dat het bewuste virus kan worden gesplitst in twee componenten, die in verwekte symptomen verschillen, doch niet in de manier van overbrengen. De scheiding in twee componenten is ons tot op heden nog niet gelukt; daarom zullen we hier blijven spreken van één virus en wel van het bloemvergroeningsvirus.

Dit virus blijkt allerm minst specifiek te zijn voor één bepaalde plantesoort; FRAZIER & POSNETTE (1956, 1957) en EVENHUIS (1957) vermelden een groot aantal waardplanten, die tot zeer verschillende families behoren. Er blijken eveneens verschillende vectoren te bestaan. FRAZIER & POSNETTE (1956, 1957) vermelden *Euscelis lineolatus* BRULLÉ, *E. plebejus* (FALL.) en *Macrosteles viridigriseus* (EDWARDS) als zodanig, BOVEY (1957) en EVENHUIS (1957) ook nog *Aphrodes bicinctus* (SCHRANK).

In 1957 werden nog enkele overbrengingsproeven met verschillende andere soorten cicaden verricht. Zo werd met *Philaenus spumarius* (L.) en *Agallia consobrina* (CURT.), die beide polyfaag zijn, een groot aantal proeven opgezet; in geen enkel geval werd echter overbrenging van het virus verkregen. Wel bleek het virus te kunnen worden overgebracht door *Aphrodes albifrons* (L.) (in twee gevallen) en *Macrosteles cristatus* (RIBAUT) (in één geval). Dat er slechts weinig gevallen van overbrenging met deze beide soorten werden verkregen, is het gevolg van het nog geringe aantal proeven, dat met deze soorten werd uitgevoerd. Bovendien bedraagt de duur van de latente periode van het virus in het insect in de proeven in de warme kas meestal zes tot zeven weken en is het vaak moeilijk – vooral bij de tere soorten – om de dieren zo lang in leven te houden. De kans is groot, dat bij verder onderzoek zal blijken, dat nog meer soorten cicaden het bloemvergroeningsvirus kunnen overbrengen; vooral in de geslachten *Macrosteles* en *Aphrodes* is deze kans groot. Op het ogenblik zijn er dus zes verschillende vectoren van het virus bekend.

Niet alle vectoren zijn even effectief, zo blijkt *Aphrodes bicinctus* een veel minder effectieve overbrenger te zijn dan *Euscelis plebejus* en *E. lineolatus*. Trouwens ook alle waardplanten zijn niet even geschikt; zo gelukte de overbrenging van het virus op klaver (witte, rode en inkarnaatklaver) veel beter dan op aster.

¹⁾ Aangenomen voor publikatie 20 mei 1958.

SUMMARY

In literature the leafhopper species *Euscelis plebejus* (FALL.), *E. lineolatus* BRULLÉ, *Macrosteles viridigriseus* (EDWARDS) and *Aphrodes bicinctus* (SCHRANK) are mentioned as vectors of the virus causing phyllody in clover and other flowers. It was also found that *Macrosteles cristatus* (RIBAUT) and *Aphrodes albifrons* (L.) are able to transmit the virus. It seems probable that more vectors of the virus exist. However in many experiments with *Philaenus spumarius* (L.) and *Agallia consobrina* (CURT.) no transmission of the virus was obtained.

LITERATUUR

- Bos, L., - 1957. Plant teratology and plant pathology. T. Pl.ziekten 63:222-231.
BOVEY, R., - 1957. Une anomalie des fleurs du trèfle causé par un virus transmis par des cicadelles. Rev. rom. Agric. Vitic. Arboric. 13:106-108.
EVENHUIS, H. H., - 1958. Investigations on a leafhopper-borne clover virus. Proc. 3rd Conf. Potato Virus Diseases, Lisse-Wageningen, 1957: 251-254.
FRAZIER, N. W. & A. F. POSNETTE, - 1956. Leafhopper transmission of a clover virus causing green petal disease in strawberry. Nature 177:1040-1041.
FRAZIER, N. W. & A. F. POSNETTE, - 1957. Transmission and host-range studies of strawberry green-petal virus. Ann. appl. Biol. 45:580-588.

Mededelingen van het Instituut voor Plantenziektenkundig Onderzoek

Contributions of the Institute for Phytopathological Research

Binnenhaven 4a, Wageningen, The Netherlands

- No 137. KLINKENBERG, C. H. en J. W. SEINHORST, De nematicide werking van Na N-methyl dithiocarbamaat (Vapam) bij toepassing in de herfst. (The nematicidal properties of Na N-methyl dithiocarbamate (Vapam) when applied in autumn). Meded. L.H.S. Gent, XXI: 397-400, 1956. Prijs no 137 en 138 samen f 0,40.
- No 138. SEINHORST, J. W., J. D. BIJLOO en C. H. KLINKENBERG, Een vergelijking van de nematicide werking van DD en van 3-5-dimethyltetrahydro-1-3-5-2H-thiadiazine-2-thion. (A comparison of the nematicidal properties of DD and of 3-5-dimethyltetrahydro-1-3-5-2H-thiadiazine-2-thione). Meded. L.H.S. Gent, XXI: 387-395, 1956. Prijs no 137 en 138 samen f 0,40.
- No 139. HOOF, H. A. VAN en SJ. TOLMA, Virusziekte bij rabarber. (Virus disease of rhubarb). Meded. Dir. Tuinb., 19: 761-764, 1956. Prijs no 139 en 140 samen f 0,60.
- No 140. HOOF, H. A. VAN, Verschil in reactie van wilde sla ten opzichte van besmetting met het slamozaiekvirus. (Differences in reaction of wild lettuces with regard to infection by *Lactuca virus 1*). Tijdschr. o. Plantenz., 62: 285-290, 1956. Prijs no 139 en 140 samen f 0,60.
- No 141. BEEMSTER, A. B. R., Onderzoekingen over een virusziekte bij stoppelknollen (*Brassica rapa* var. *rapifera*). (Investigations on a virus disease of turnip). Tijdschr. o. Plantenz., 63: 1-12, 1957. Prijs f 0,40.
- No 142. FRANSSEN, C. J. H., De levenswijze en de bestrijding van de tuinboonkever. (The biology and control of *Bruchus rufimanus* Boh.). Versl. Landbouwk. Onderz., No. 62. 10, 75 pp., 1956. Prijs f 2,75.
- No 143. FRANSSEN, C. J. H., De levenswijze en de bestrijding van de bonekever (*Acanthoscelides obtectus* Say). (Biology and control of the common bean weevil). Meded. Dir. Tuinb., 19: 797-809, 1956. Prijs f 0,55.
- No 144. NUVELDT, W. C., Levenswijze en bestrijding van de aspergevlug. (*Platyparea poeciloptera* Schrank) in Nederland. (Biology and control of *Platyparea poeciloptera* Schrank in the Netherlands). Versl. Landbouwk. Onderz., No 63. 4, 40 pp., 1956. Prijs f 1,65.
- No 145. VAUGHAN, EDWARD K., A device for the rapid removal of tannins from virus infected plant tissues before extraction of inoculum. Tijdschr. o. Plantenz., 266-270, 1956. Prijs no 145 en 146 samen f 0,30.
- No 146. VAUGHAN, EDWARD K., Attempts to transfer *Rubus* and *Fragaria* viruses into herbaceous hosts. Tijdschr. o. Plantenz., 62: 271-273, 1956. Prijs no 145 en 146 samen f 0,30.
- No 147. ZEYLSTRA, H. H., Papierchromatografie als middel voor de diagnose van de ringvlekkenziekte van zoete kers. Een voorlopige mededeling. (Paper chromatography and diagnosis of ring spot disease in sweet cherry. A preliminary report). Tijdschr. o. Plantenz., 62: 325-326, 1956. Prijs no 147 en 148 samen f 0,25.
- No 148. QUAK, FRED. a., Meristeencultuur, gecombineerd met warmtebehandeling, voor het verkrijgen van virusvrije anjerplanten. (Meristem culture, combined with heat treatment, in order to obtain virus-free carnation plants). Tijdschr. o. Plantenz., 63: 13-14, 1957. Prijs no 147 en 148 samen f 0,25.
- No 149. FRANSSEN, J. J., De landbouwluchtvaart in Nederland. (Agricultural aviation in the Netherlands). Landbouwvoorl., 13: 578-585, 1956. f 0,45.
- No 150. HUBBELING, N., New aspects of breeding for disease resistance in beans. *Euphytica*, 6: 111-141, 1957. Prijs no 150 f 3,-.
- No 151. QUAK, FRED. a., Bladvlekkenziekte bij spruitkool veroorzaakt door *Mycosphaerella brassicicola* (Fr.) Lindau. (Ringspot disease of brussels sprouts caused by *Mycosphaerella brassicicola* (Fr.) Lindau) Meded. Dir. Tuinb., 20: 317-320, 1957. Prijs no 151, 152 en 153 samen f 0,95.
- No 152. LABRUYÈRE, R. E., Enkele waarnemingen over de schimmel *Elsinoë veneta* (Burkh.) Jenk., de perfecte vorm van *Sphaceloma necator* (Ell. et Ev.) Jenk. en Shear. (Observations on *Elsinoë veneta* (Burkh.) the perfect form of *Sphaceloma necator* (Ell. et Ev.) on raspberry) Tijdschr. o. Plantenz., 63: 153-158, 1957. Prijs no 151, 152 en 153 samen f 0,95.

- No 176. s'JACOB, J. C., Phytopathologische Probleme bei der Resistenzzüchtung. *Z. Pfl.-zücht.*, 40: 113-124, 1958. Prijs no 176 f 0,50.
- No 177. HOOF, H. A. VAN, Some considerations on the transmission of non-persistent viruses by aphids. *Proc. 3rd Conf. Potato Virus Diseases*: 114-116, 1958. Prijs no 177 f 0,25.
- No 178. EVENHUIS, H. H., Investigations on a leafhopper-borne clover virus. *Proc. 3rd Conf. Potato Virus Diseases*: 251-254, 1958. Prijs no 178 en 179 samen f 0,25.
- No 179. EVENHUIS, H. H., De vectoren van het bloemvergroeningsvirus van klaver. (The vectors of the virus causing phyllody (virescence) in clover flowers) *T. Pl.-ziekten*, 64: 335-336, 1958. Prijs no 178 en 179 samen f 0,25.
- No 180. BEEMSTER, A. B. R., Some aspects of mature plant resistance to viruses in the potato. *Proc. 3rd Conf. Potato Virus Diseases*: 212-217, 1958. Prijs no 180 f 0,35.
- No 181. EVENHUIS, H. H., Over de invloed van de winter op de parasitering van de appelbloedluis, *Eriosoma lanigerum*, door haar parasiet *Aphelinus mali*. (The influence of the winter on the parasitism of the woolly aphid, *Eriosoma lanigerum*, by its parasite *Aphelinus mali*) *T. Pl.-ziekten*, 64: 328-332, 1958. Prijs no 181 f 0,30.
- No 182. BOS, L. en J. P. H. VAN DER WANT, Virusziekten van vlinderbloemigen. (Virus diseases of leguminous plants) *Landbouwvoorlichting*, 15: 550-558 en 573-587, 1958. Prijs no 182 f 1,15.
- No 183. Franssen, C. J. H. en P. Huisman, De levenswijze en bestrijdingsmogelijkheden van de vroege akkertrips (*Thrips angusticeps* Uzel. (The biology and control of *Thrips angusticeps* Uzel) *Versl. landbouwk. Onderz.*, 64. 10: 1-104, 1958. Prijs no 183 f 3,50
- No 184. Fluiter, H. J. de, Bladluisbestrijding ter voorkoming van virusverspreiding in aardbeien. (Aphid control in strawberries to prevent spread of strawberry viruses) *Meded. Landbouwhoges. Gent*, 23:745-769, 1958. Prijs no 184 f 0,95